Entanglement molecules

W. Dür
Institut für Theoretische Physik, Universität Innsbruck, A-6020 Innsbruck, Austria
(February 1, 2008)

arXiv:quant-ph/0006105v1 22 Jun 2000

We investigate the entanglement properties of multiparticle systems, concentrating on the case where the entanglement is robust against disposal of particles. Two qubits —belonging to a multipartite system— are entangled in this sense iff their reduced density matrix is entangled. We introduce a family of multiqubit states, for which one can choose for any pair of qubits independently whether they should be entangled or not as well as the relative strength of the entanglement, thus providing the possibility to construct all kinds of "Entanglement molecules". For some particular configurations, we also give the maximal amount of entanglement achievable.

03.67.-a, 03.65.Bz, 03.65.Ca, 03.67.Hk

Entanglement is at the heart of Quantum Information theory. In recent years, there has been an ongoing effort to characterice quantitatively and qualitatively entanglenglement. While for bipartite systems this problem is essentially solved, it remains still open for multipartite systems. In this case, there exist several possible approaches to identify different kinds of multipartite entanglement, and many interesting phenomena related to multipartite entanglement have been discovered [1].

Here we concentrate on bipartite aspects of multipartite entanglement, in particular on bipartite entanglement which is robust against disposal of particles. We consider N spatially separated parties $A_1, \ldots, A_N$, each possessing a qubit. We first investigate the N–party Greenberger-Horne-Zeilinger (GHZ)-state [2]

$$|GHZ\rangle = \frac{1}{\sqrt{2}}(|0^{\otimes N}\rangle + |1^{\otimes N}\rangle), \tag{1}$$

which is considered to be a maximally entangled state (MES) of N particles in several senses. For example, one can create a MES shared by any two of the parties with help of a (local) measurement performed by the remaining ones. Thus any two particles are potentially entangled, i.e. when allowing for assistance of the other parties, bipartite entanglement can be obtained from state $|GHZ\rangle$. However, it is essential that the remaining $(N-2)$ parties perform measurements to assist the other two parties to share entanglement. If however —for some reason— the information about only one of the particles, say A_N, is lost (or party A_N decides not to cooperate with the remaining ones), the state of the remaining parties is only classically correlated and thus not entangled. In particular, the reduced density operator of any two parties is separable[1]. When considering the reduced density operator of two parties, we deal with the situation where the information about all remaining particles is not accessable (or the remaining parties are not willing to cooperate).

We thus say that two particles are entangled iff their reduced density operator is non–separable, i.e. the two particles share entanglement, independent what happens to the remaining particles. Such a definition is very suitable from a practical point of view, as there are certain multipartite scenarious where one is interested in entanglement properties of pairs of parties, which are independent of other parties. Note that in this sense, the state $|GHZ\rangle$ contains no (bipartite) entanglement at all.

But are there N–particle states which are still entangled when tracing out *any* $(N-2)$ particles, i.e. are there states where all particles are entangled with all other particles ? And if this is possible, what is the maximal amount of entanglement the remaining two parties can share ? In this paper, we will answer these questions and we will consider the more general setup where some parties are entangled, while some others are not. For example, for $N = 3$ one may have that the reduced density operators ρ_{12} and ρ_{13} remain entangled, but ρ_{23} is separable. We will show that one can have all possible configurations of this kind, i.e. there exist states where one can choose for each of the reduced density operators ρ_{kl}, $k < l \in \{1, \ldots, N\}$ independently whether it should be entangled or not. This allows to build general structures of N particle states, which we call 'Entanglement molecules' in spirit of the generalization of Wootters idea of an 'Entangled chain' [3], where one has a string of qubits, each qubit being entangled only with its nearest neighbours. We are considering more general setups, e.g. closed rings of particles where one only has nearest neighbour entanglement, (finite) strings with distance dependent entanglement (also 2nd and 3rd and so on neighbourhood entanglement), entanglement buckiballs (like the C_{60} molecule), or more generally all possible setups of this kind one may imagine.

There are $N(N-1)/2$ different bipartite reduced density operators ρ_{kl} which may be either separable or not. If the reduced density operator ρ_{kl} is non–separable, this

[1] Given a N–partite state ρ, the reduced density operator ρ_{12} of party A_1 and A_2 is defined as $\rho_{12} \equiv \mathrm{tr}_{3,\ldots,N}(\rho)$. The operator ρ_{12} is separable if it can be written as a convex combination of product states.

automatically implies that a MES shared between parties A_k and A_l can be distilled (when allowing for several copies of the state), even without help of the remaining parties. This is due to the fact that for two qubit systems, inseparability is equivalent to distillability [4]. In fact, the remaining parties can by no means prevent parties A_k and A_l to distill a MES. In a diagram, this will be visualized by a line between particles A_k and A_l which represents entanglement between the two parties in question (see Fig. 1). Each of the particles may be entangled with one (or more) of the remaining $(N-1)$ particles. In particular one can have that any particle is entangled with all the remaining ones. Clearly, it is interesting to ask how strong these 'bindings' (the entanglement between two particles) can be. Therefore, one has to quantify the entanglement of the bipartite reduced density operators ρ_{kl}. In this work, we choose as a measure of entanglement the concurrence $\mathcal{C}$ (for a definition of the concurrence see e.g. [5]). On one hand, we follow the lines suggested in [3,5], on the other we have that the entanglement of formation —the amount of entanglement required to prepare a state ρ— is monotonically increasing with the concurrence $\mathcal{C}$ [2] and thus the concurrence itself may be used to measure the strength of the bindings. For two special cases of particular interest, we will give the states with the maximal achieveable strength of the bindings.

Let us start by introducing a family of N qubit states, which includes all possible configurations of 'Entanglement molecules'. First we specify for each of the reduced density operators ρ_{kl} whether it should be distillable or not, i.e. whether entanglement between the parties A_k and A_l can be distilled — without help of the remaining parties — or not. Let $I = \{k_1 l_1, \ldots, k_M l_M\}$ be the set of all those pairs where distillation should be possible, i.e. for $kl \in I$, we have that ρ_{kl} is distillable. We define the state

$$|\Psi_{ij}\rangle \equiv |\Psi^+\rangle_{ij} \otimes |0\ldots 0\rangle_{\text{rest}}, \tag{2}$$

that is the particles A_i and A_j are in a MES, namely $|\Psi^+\rangle = 1/\sqrt{2}(|01\rangle + |10\rangle)$, and the remaining particles are disentangled from each other and from $A_i A_j$. We now introduce a family of states which has the desired properties:

$$\rho_I = \frac{1}{M} \sum_{kl \in I} x_{kl} |\Psi_{kl}\rangle\langle\Psi_{kl}|, \tag{3}$$

where $M \equiv \sum_{kl\in I} x_{kl}$ is a normalization factor. It is straightforward to calculate the reduced density operators ρ_{kl}. In the standard basis, one can check for $kl \in I$ that ρ_{kl} is of the form

$$\rho_{kl} = \frac{1}{M} \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & b & x_{kl}/2 & 0 \\ 0 & x_{kl}/2 & c & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{4}$$

while for $mn \notin I$ one obtains

$$\rho_{mn} = \frac{1}{M} \begin{pmatrix} \tilde{a} & 0 & 0 & 0 \\ 0 & \tilde{b} & 0 & 0 \\ 0 & 0 & \tilde{c} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{5}$$

It is simple to calculate the concurrence in both cases:

$$\begin{aligned} \mathcal{C}_{kl} &= \frac{x_{kl}}{M} \text{ iff } kl \in I \\ \mathcal{C}_{mn} &= 0 \text{ iff } mn \notin I, \end{aligned} \tag{6}$$

where we denote $\mathcal{C}_{kl} \equiv \mathcal{C}(\rho_{kl})$. We have that in $\mathbb{C}^2 \otimes \mathbb{C}^2$ systems, nonzero concurrence automatically implies distillability of the state, while zero concurrence implies separability. Thus it is already clear that ρ_I has the desired properties, i.e. entanglement between A_k and A_l can be distilled iff $kl \in I$. We see that we can arbitrarily choose the relative strength of the bindings (measured by the concurrence) via the positive coefficients x_{kl}. It is now straightforward to explicitly construct the examples illustrated in Fig. 1: In (a) we have that $x_{12} = x_{34} = x_{56} = 2/9, x_{23} = x_{45} = x_{16} = 1/9$ (entanglement ring); In (b) $x_{kl} = 1/6$ iff both k and l are even [odd] respectively and zero otherwise (all even [odd] particles are equally entangled) ; In (c) we have that $x_{1l} = 1/5$, while all other $x_{kl} = 0$ (one particle equally entangled with five other ones); Finally, in (d) $x_{kl} = 1/15$ (all particles equally entangled). In a similar way, one can construct all examples mentioned in the introduction, such as strings with distance dependent entanglement or entanglement buckiballs.

Note that in this construction, we have that

$$\sum_{i<k} \mathcal{C}_{ik} = 1, \tag{7}$$

which shows that in a situation where all bindings have the same strength, the concurrence of the corresponding reduced density operators is determined by the total number of bindings, i.e. the number of elements in I. Hence, a state constructed in this way, where each particle is entangled with all others (i.e. all possible reduced density operators are distillable) and the strength of all bindings is equal has $\mathcal{C}_{kl} = 2/[N(N-1)]$. As we shall see next, this is however not the maximum value one can achieve for this particular configuration. In the following, we will analyze the two situations illustrated in Fig. 1(c) and (d), namely —(i) all particles are equally entangled and (ii) one particle which is equally entangled with $(N-1)$ others— and determine the maximum strength of the bindings for $N = 3$.

(i) All particles pairwise entangled

We start with the case where all particles are equally

[2] The entanglement of formation is given by $E_f(\rho) = h(\frac{1}{2} + \frac{1}{2}\sqrt{1-\mathcal{C}^2})$, where $\mathcal{C}$ is the concurrence and h is the binary entropy function $h(x) = -x\log_2 x - (1-x)\log_2(1-x)$.

entangled with each other (see also Fig. 1(d)). As shown in [6], the state

$$|W\rangle = 1/\sqrt{3}(|001\rangle + |010\rangle + |100\rangle), \tag{8}$$

is the state of three qubits whose entanglement has the highest degree of endurance against loss of one of the three qubits. In particular, $|W\rangle$ maximizes the function

$$\mathcal{E}_{\min}(\psi) \equiv \min(\mathcal{C}_{12}, \mathcal{C}_{13}, \mathcal{C}_{23}), \tag{9}$$

and has $\mathcal{C}_W \equiv \mathcal{C}_{12} = \mathcal{C}_{13} = \mathcal{C}_{23} = 2/3$. It follows that $|W\rangle$ is the 3–qubit state where all particles are equally entangled and the entanglement — measured by the concurrence $\mathcal{C}$— is maximal. Comparing $|W\rangle$ with ρ_I of the form (3) with $I = \{12, 13, 23\}$, one immeadetly observes that $\mathcal{C}_{\rho_I} = 1/3$, while $\mathcal{C}_W = 2/3$.

More generally, let us consider the N–party form $|W_N\rangle$ of the state $|W\rangle$, defined as

$$|W_N\rangle \equiv 1/\sqrt{N}|N-1, 1\rangle, \tag{10}$$

where $|N-1, 1\rangle$ denotes the (unnormalized) totally symmetric state including $N-1$ zeros and 1 ones, e.g. $|2, 1\rangle = |001\rangle + |010\rangle + |100\rangle$. As shown in [6], state $|W_N\rangle$ is a N qubit state with all reduced density operators equal and the concurrences are given by $\mathcal{C}_{kl} = 2/N$, which has to be compared to $\mathcal{C}_{\rho_I} = 2/[N(N-1)]$, ρ_I being a state of the family (3) where all particles are equally entangled. It is however not known whether $\mathcal{C}_{kl} = 2/N$, $\forall k \neq l$ is the maximal value achievable.

(ii) One particle entangled with $N-1$ others We consider now the case where one particle is equally entangled with $(N-1)$ others and determine the maximal possible strength $\mathcal{C}$ of the bindings (see Fig. 1(c)). A related problem of this kind, namely the question of optimal entanglement splitting, i.e. the optimal way for a party B to equally distribute its initial entanglement (shared with a party A) among several partners, was recently analyzed by Bruß in [7].

As shown in [5], we have for $N = 3$ that $\mathcal{C}_{12}^2 + \mathcal{C}_{13}^2 \leq 1$, i.e for $\mathcal{C}_{12} = \mathcal{C}_{13} \equiv \mathcal{C}$ we have that $\mathcal{C} \leq 1/\sqrt{2}$. This value is achieved by the state

$$|\psi\rangle = \frac{1}{\sqrt{2}}|100\rangle + \frac{1}{2}(|001\rangle + |010\rangle). \tag{11}$$

More generally, it was conjectured in [5] that the inequality $\sum_{k=2}^{N} \mathcal{C}_{1k}^2 \leq 1$ also holds. If we demand again that $\mathcal{C}_{1k} = \mathcal{C}_{1l} \equiv \mathcal{C}$ $\forall k, l > 1$, we find that $\mathcal{C} \leq 1/\sqrt{N-1}$. This value is obtained by the state

$$|\psi\rangle = a|1\rangle|00\ldots0\rangle + b|0\rangle|N-2, 1\rangle), \tag{12}$$

with $a = 1/\sqrt{2}$ and $b = 1/\sqrt{2(N-1)}$, where $|N-2, 1\rangle$ is again an (unnormalized) totally symmetric state including $(N-2)$ zeros and one 1. For a state of the form (3), where one particle is equally entangled with $(N-1)$ others, one obtains $\mathcal{C}_{1k} = 1/(N-1)$.

For practical purposes, it may sometimes be useful to consider the fidelity of the reduced density operator (i.e. the maximal overlap of the reduced density operator with any MES) —which is generally not a proper measure of entanglement— instead of the concurrence. The fidelity of the reduced density operator however indicates the achievable quality to perform certain quantum information tasks, e.g. teleportation [8]. The fidelity of a density operator ρ is defined as

$$F_\rho = \max\langle\Phi|\rho|\Phi\rangle, \tag{13}$$

where the maximum is taken over all maximally entangled states $|\Phi\rangle$. In the situation we consider, where one particle is equally entangled (i.e. F_ρ is equal for all reduced density operators ρ_{1k}) with $(N-1)$ others, one can derive a bound for the maximum fidelity F_ρ of the reduced density operators ρ_{1k} using results from optimal cloning [9]. Given a density operator ρ with a certain fidelity F_ρ, one can use ρ to teleport [8] the (unknown) state of a particle. As shown by the Horodecki [10], one can find a teleportation protocol which works equally well for all input states and has the maximum teleport fidelity $F_t = (2F_\rho + 1)/3$. In our situation, we have that particle A_1 is entangled with $(N-1)$ other particles. Thus, when performing a certain teleportation protocol [11,12], one obtains $(N-1)$ (imperfect) clones of a state at the locations $A_2, \ldots, A_N$, where the quality of the clones is determined by the teleport fidelity F_t of the corresponding reduced density operators. As shown by Werner [9], the maximum cloning fidelity of a $1 \to (N-1)$ cloner is given by $F_c = [2(N-1)+1]/[3(N-1)]$, from which follows that the teleport fidelity F_t of the reduced density operators must fulfill $F_t \leq F_c$, otherwise one could construct in this way a cloning machine which works better than the optimal one, which is clearly impossible. We thus have that

$$F_\rho \leq \frac{1}{2} + \frac{1}{2(N-1)}. \tag{14}$$

For the state $|\psi\rangle$ which maximizes the concurrence, we find that it does not obtain the maximal possible fidelity F_ρ. However, there exist states for which the maximal possible value of the fidelity is obtained. For example, a state of the form (12) with $a = 1/\sqrt{N(N-1)}$ and $b = \sqrt{(N-1)/N}$ has the desired properties, as well as the 'telecloning state' introduced in [11]. On the other hand, a state of the form (3), where one particle is equally entangled with $(N-1)$ others has $F_\rho = [2+N]/[4(N-1)]$, much smaller than the optimal value.

Note that the maximum value of the entanglement (measured by the concurrence) of a given qubit with its neighbours is not simply determined by the number of entangled neighbours, but also by the properties of the neighbours (i.e. the number of particles to which the neighbouring particles are entangled). This can be seen by noting that (i) in the case where one particle is equally entangled with two others (which are disentangled among

themselves), the maximimal value for the concurrence is given by $\mathcal{C} = 1/\sqrt{2}$ and (ii) in the case where three qubits are equally entangled, the maximum value is given by $\mathcal{C} = 2/3$, which shows that the entanglement between systems A_2 and A_3 influence the maximum value of the entanglement between systems A_1A_2 and A_1A_3.

In summary, we provided a family of states which allows to construct all possible kinds of 'Entanglement molecules', i.e. one can choose for any pair of qubits independently whether a MES can be distilled without help of the remaining parties or not. In addition, the relative strength of the bindings (measured by the concurrence) can be adjusted arbitrarily. We investigated two particular configurations closer and provided states achieving the maximum value for the strength of the bindings.

I would like to thank G. Vidal and J. I. Cirac for discussions. This work was supported by the Austrian Science Foundation under the SFB "control and measurement of coherent quantum systems (Project 11), the European Community under the TMR network ERB–FMRX–CT96–0087 and project EQUIP (contract IST-1999-11053), the European Science Foundation, and the Institute for Quantum Information GmbH.



FIG. 1. Several kinds of Entanglement molecules for $N = 6$. (a) Entanglement ring - double lines indicate stronger entanglement between particles. (b) All even [odd] particles are entangled respectively. (c) One particle A_1 equally entangled with five others. (d) All particles equally entangled.

[1] A. Thapliyal, Phys. Rev. A **59** (1999) 3336; J. Kempe, Phys. Rev. A **60** (1999) 910; C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, Phys. Rev. Lett. **83**, 3081 (1999); W. Dür, J.I. Cirac and R. Tarrach, Phys. Rev. Lett. **83** (1999) 3562; D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, B. M. Terhal, quant-ph/9908070; J. A. Smolin, quant-ph/0001001; W. Dür and J. I. Cirac, quant-ph/0002028 (to appear in Phys. Rev. A); P. W. Shor, J. A. Smolin and A. V. Thapliyal, quant-ph 0005117.

[2] D. M. Greenberger, M. Horne, A. Zeilinger, *Bell's theorem, Quantum Theory, and Conceptions of the Universe,* ed. M. Kafatos, Kluwer, Dordrecht 69 (1989);

[3] W. K. Wootters, quant-ph/0001114

[4] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. Lett. **78**, 574 (1997).

[5] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A **61**, 052306 (2000); see also quant-ph/9907047

[6] W. Dür, G. Vidal and J. I. Cirac, quant-ph/0005115

[7] D. Bruß, quant-ph/9902023

[8] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett **70**, 1895-1899 (1993)

[9] R. F. Werner, quant-ph/9804001

[10] P. Horodecki, M. Horodecki and R. Horodecki, quant-ph/9807091

[11] M. Murao, D. Jonathan, M. B. Plenio and V. Vedral, Phys. Rev. A. **59**, 156 (1999)

[12] W. Dür and J. I. Cirac, Journal of Modern Optics, 2000, Vol. 47, No. 2/3, 247-255